

The quasi-symmetric Hölder equivalence problem


P. Pansu

August 27th, 2015

Today's menu: a problem in metric geometry motivated by negative curvature.

Main actor: complex hyperbolic plane $H_{\mathbb{C}}^2$.

The ball $B \subset \mathbb{C}^2$ has biholomorphism group $PU(2, 1)$, which preserves a complete Riemannian metric on B and the contact structure of complex tangents on ∂B . $PU(2, 1)$ has a subgroup $S = \mathbb{R} \times Heis$ acting simply transitively on the ball. The conjugation action of \mathbb{R} on $Heis$ is by dilations, $Lie(Heis) = V_1 \oplus V_2$, $\delta_t = e^{it/4}$ on V_j . The induced metric on S is of the form $dt^2 + \delta_t^* g_0 = dt^2 + e^{t/2} g_{V_1} + e^t g_{V_2}$, for some left-invariant Riemannian metric g_0 on $Heis$. \mathbb{R} factors are geodesics, $Heis$ -orbits are horospheres. $\frac{\partial}{\partial t}$ and V_2 generate a complex geodesic, curvature -1 . Each line in V_1 generates with $\frac{\partial}{\partial t}$ a Lagrangian plane, curvature $-\frac{1}{4}$.



Definition

Let M be a Riemannian manifold. Let $-1 \leq \delta < 0$. Say M is δ -pinched if sectional curvature ranges between -1 and δ . Define the optimal pinching $\delta(M)$ of M as the least $\delta \geq -1$ such that M is quasi-isometric to a δ -pinched complete simply connected Riemannian manifold.

Definition

Let M be a Riemannian manifold. Let $-1 \leq \delta < 0$. Say M is δ -pinched if sectional curvature ranges between -1 and δ . Define the optimal pinching $\delta(M)$ of M as the least $\delta \geq -1$ such that M is quasi-isometric to a δ -pinched complete simply connected Riemannian manifold.

Example: Complex hyperbolic plane $H_{\mathbb{C}}^2$ is $-\frac{1}{4}$ -pinched. Is it true that $\delta(H_{\mathbb{C}}^2) = -\frac{1}{4}$?


Definition

Let M be a Riemannian manifold. Let $-1 \leq \delta < 0$. Say M is δ -pinched if sectional curvature ranges between -1 and δ . Define the optimal pinching $\delta(M)$ of M as the least $\delta \geq -1$ such that M is quasi-isometric to a δ -pinched complete simply connected Riemannian manifold.

Example: Complex hyperbolic plane $H_{\mathbb{C}}^2$ is $-\frac{1}{4}$ -pinched. Is it true that $\delta(H_{\mathbb{C}}^2) = -\frac{1}{4}$?

Facts.

- Negatively curved manifolds M have a visual sphere ∂M , equipped with a visual metric.
- If M is δ -pinched, polar coordinates define a $C^{\sqrt{-\delta}}$ -Hölder homeomorphism from the round sphere $S \rightarrow \partial M$, with 1-Lipschitz inverse.
- Quasi-isometries between negatively curved Riemannian manifolds induce *quasisymmetric* maps between ideal boundaries.





Definition

Let M be a Riemannian manifold. Let $-1 \leq \delta < 0$. Say M is δ -pinched if sectional curvature ranges between -1 and δ . Define the optimal pinching $\delta(M)$ of M as the least $\delta \geq -1$ such that M is quasi-isometric to a δ -pinched complete simply connected Riemannian manifold.

Example: Complex hyperbolic plane $H_{\mathbb{C}}^2$ is $-\frac{1}{4}$ -pinched. Is it true that $\delta(H_{\mathbb{C}}^2) = -\frac{1}{4}$?

Facts.

- Negatively curved manifolds M have a visual sphere ∂M , equipped with a visual metric.
- If M is δ -pinched, polar coordinates define a $C^{\sqrt{-\delta}}$ -Hölder homeomorphism from the round sphere $S \rightarrow \partial M$, with 1-Lipschitz inverse.
- Quasi-isometries between negatively curved Riemannian manifolds induce *quasisymmetric* maps between ideal boundaries.



Definition

$\alpha_{qs}(X) = \sup\{\alpha \in (0, 1) \mid \exists \text{ locally a } C^\alpha \text{ homeomorphism with Lipschitz inverse from Euclidean space to a metric space quasisymmetric to } X\}$.

By definition, $\sqrt{-\delta(M)} \leq \alpha_{qs}(\partial M)$.

Example: The visual boundary of complex hyperbolic plane is a sub-Riemannian 3-sphere, quasisymmetric to *Heis*. Note that $\alpha_{qs}(\text{Heis}) \geq \frac{1}{2}$.

Definition

$\alpha_{qs}(X) = \sup\{\alpha \in (0, 1) \mid \exists \text{ locally a } C^\alpha \text{ homeomorphism with Lipschitz inverse from Euclidean space to a metric space quasisymmetric to } X\}$.

By definition, $\sqrt{-\delta(M)} \leq \alpha_{qs}(\partial M)$.

Example: The visual boundary of complex hyperbolic plane is a sub-Riemannian 3-sphere, quasisymmetric to *Heis*. Note that $\alpha_{qs}(\textit{Heis}) \geq \frac{1}{2}$.

Conjecture

$$\alpha_{qs}(\textit{Heis}) = \frac{1}{2}.$$

I will explain:

Theorem

$$\alpha_{qs}(\textit{Heis}) \leq \frac{2}{3}.$$

Note that consequence $\delta(H_{\mathbb{C}}^2) \geq -\frac{4}{9}$ was already known.

Question (Hölder equivalence problem, Gromov 1993)

Let X be a sub-Riemannian manifold. For which $\alpha \in (0, 1)$ does there exist locally a homeomorphism from Euclidean space to X which is C^α -Hölder continuous ?

Question (Hölder equivalence problem, Gromov 1993)

Let X be a sub-Riemannian manifold. For which $\alpha \in (0, 1)$ does there exist locally a homeomorphism from Euclidean space to X which is C^α -Hölder continuous ?

Definition

Let $\alpha(X) = \sup\{\alpha \in (0, 1) \mid \exists \text{ locally a } C^\alpha \text{ homeomorphism } \mathbb{R}^n \rightarrow X\}$.

Question (Hölder equivalence problem, Gromov 1993)

Let X be a sub-Riemannian manifold. For which $\alpha \in (0, 1)$ does there exist locally a homeomorphism from Euclidean space to X which is C^α -Hölder continuous ?

Definition

Let $\alpha(X) = \sup\{\alpha \in (0, 1) \mid \exists \text{ locally a } C^\alpha \text{ homeomorphism } \mathbb{R}^n \rightarrow X\}$.

Theorem (Gromov 1993)

Let metric space X have dimension n , Hausdorff dimension Q . Then $\alpha(X) \leq \frac{n}{Q}$.

Let sub-Riem. X have dimension n , Hausdorff dimension Q . Then $\alpha(X) \leq \frac{n-1}{Q-1}$.

Let X be a $2m + 1$ -dimensional contact manifold. Then $\alpha(X) \leq \frac{m+1}{m+2} (\leq \frac{2m}{2m+1})$.

Best known bound $\alpha(\text{Heis}) \leq \frac{2}{3}$.

Gromov's slogan:

Here
Euclidean
\exists k -dimensional subset with Hausdorff dimension $\leq k$

There
Sub-Riemannian
\forall k -dimensional subset, Hausdorff dimension $\geq k'$

then $\alpha(\text{Sub-Riemannian}) \leq \frac{k}{k'}$.

Gromov's slogan:

Here
Euclidean
$\exists k$ -dimensional subset with Hausdorff dimension $\leq k$

There
Sub-Riemannian
$\forall k$ -dimensional subset, Hausdorff dimension $\geq k'$

then $\alpha(\text{Sub-Riemannian}) \leq \frac{k}{k'}$.

In my argument,

Here
Euclidean
explicit family of subsets with Hausdorff dimension $\leq k$

There
Heisenberg
for almost every image subset, Hausdorff dimension $\geq k'$

then $\alpha(\text{Heisenberg}) \leq \frac{k}{k'}$.

Gromov's slogan:

Here
Euclidean
\exists k -dimensional subset with Hausdorff dimension $\leq k$

There
Sub-Riemannian
\forall k -dimensional subset, Hausdorff dimension $\geq k'$

$$\text{then } \alpha(\text{Sub-Riemannian}) \leq \frac{k}{k'}.$$

In my argument,

Here
Euclidean
explicit family of subsets with Hausdorff dimension $\leq k$

There
Heisenberg
for almost every image subset, Hausdorff dimension $\geq k'$

$$\text{then } \alpha(\text{Heisenberg}) \leq \frac{k}{k'}.$$

Furthermore, only quasisymmetric invariants are used, whence

$$\alpha_{qs}(\text{Heisenberg}) \leq \frac{k}{k'}.$$

Let $f : \mathbb{R}^3 \rightarrow Heis$ be a Hölder homeomorphism with Lipschitz $f^{-1} : Heis \rightarrow \mathbb{R}^3$. Let $v : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a coordinate function. Then $u = v \circ f^{-1}$ is Lipschitz. Imagine that *coarea inequality* holds:

$$\int_G Lip_u^4 \leq \int_{\mathbb{R}} \left(\int_{u^{-1}(t)} Lip_u^3 \right) dt \leq \text{const.} \int_{\mathbb{R}} \mathcal{H}^3(u^{-1}(t)) dt. \quad (1)$$

Here, Lip_u denotes the local Lipschitz constant. Since, for non constant u , $\int_X Lip_u^4 > 0$, this shows that there exists $t \in \mathbb{R}$ such that $\mathcal{H}^3(u^{-1}(t)) > 0$, and therefore $u^{-1}(t)$ has Hausdorff dimension at least 3.

Let $f : \mathbb{R}^3 \rightarrow Heis$ be a Hölder homeomorphism with Lipschitz $f^{-1} : Heis \rightarrow \mathbb{R}^3$. Let $v : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a coordinate function. Then $u = v \circ f^{-1}$ is Lipschitz. Imagine that *coarea inequality* holds:

$$\int_G Lip_u^4 \leq \int_{\mathbb{R}} \left(\int_{u^{-1}(t)} Lip_u^3 \right) dt \leq \text{const.} \int_{\mathbb{R}} \mathcal{H}^3(u^{-1}(t)) dt. \quad (1)$$

Here, Lip_u denotes the local Lipschitz constant. Since, for non constant u , $\int_X Lip_u^4 > 0$, this shows that there exists $t \in \mathbb{R}$ such that $\mathcal{H}^3(u^{-1}(t)) > 0$, and therefore $u^{-1}(t)$ has Hausdorff dimension at least 3.

Unfortunately, [Magnani 2002](#)'s *coarea inequality* goes in the opposite direction!

Let $f : \mathbb{R}^3 \rightarrow Heis$ be a Hölder homeomorphism with Lipschitz $f^{-1} : Heis \rightarrow \mathbb{R}^3$. Let $v : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a coordinate function. Then $u = v \circ f^{-1}$ is Lipschitz. Imagine that *coarea inequality* holds:

$$\int_G Lip_u^4 \leq \int_{\mathbb{R}} \left(\int_{u^{-1}(t)} Lip_u^3 \right) dt \leq \text{const.} \int_{\mathbb{R}} \mathcal{H}^3(u^{-1}(t)) dt. \quad (1)$$

Here, Lip_u denotes the local Lipschitz constant. Since, for non constant u , $\int_X Lip_u^4 > 0$, this shows that there exists $t \in \mathbb{R}$ such that $\mathcal{H}^3(u^{-1}(t)) > 0$, and therefore $u^{-1}(t)$ has Hausdorff dimension at least 3.

Unfortunately, **Magnani 2002**'s *coarea inequality* goes in the opposite direction!

Strategy: replace conformally invariant integrals $\int Lip_u^4$ with *packing measures* which are quasisymmetric invariants and satisfy coarea inequality in the right direction. If possible, extend to vector valued maps u .

Let N be an integer, let $\ell \geq 1$. Let X be a metric space. An (N, ℓ) -packing is a countable collection of balls $\{B_j\}$ such that the collection of concentric balls ℓB_j has multiplicity $< N$.

Let N be an integer, let $\ell \geq 1$. Let X be a metric space. An (N, ℓ) -packing is a countable collection of balls $\{B_j\}$ such that the collection of concentric balls ℓB_j has multiplicity $< N$.

Let ϕ be a positive function on the set of balls in X . Define

$$\Phi_{N,\ell}^{p;\epsilon}(A) = \sup \left\{ \sum_i \phi(B_i)^p ; \{B_i\} (N, \ell)\text{-packing of } X, \text{ centered on } A, \text{ of mesh } \leq \epsilon \right\}.$$

Define the *packing pre-measure associated to ϕ* by

$$\Phi_{N,\ell}^p(A) = \lim_{\epsilon \rightarrow 0} \Phi_{N,\ell}^{p;\epsilon}(A).$$

Let N be an integer, let $\ell \geq 1$. Let X be a metric space. An (N, ℓ) -packing is a countable collection of balls $\{B_j\}$ such that the collection of concentric balls ℓB_j has multiplicity $< N$.

Let ϕ be a positive function on the set of balls in X . Define

$$\Phi_{N,\ell}^{p;\epsilon}(A) = \sup \left\{ \sum_i \phi(B_i)^p ; \{B_i\} (N, \ell)\text{-packing of } X, \text{ centered on } A, \text{ of mesh } \leq \epsilon \right\}.$$

Define the *packing pre-measure associated to ϕ* by

$$\Phi_{N,\ell}^p(A) = \lim_{\epsilon \rightarrow 0} \Phi_{N,\ell}^{p;\epsilon}(A).$$

Example: $\pi(B) = \text{radius}(B)$ leads to “usual” packing measure Π^p and packing dimension. It is Hölder covariant.

Example: let $u : X \rightarrow M$ measure space. $e_u(B) = \text{measure}(u(B))$ leads to p -energy E_u^p . It is quasisymmetry invariant: if g is quasisymmetric, $\forall \ell, \exists \ell'$ such that

$$E_{u,N,\ell}^p(A) \leq E_{u',N,\ell'}^p(g(A)).$$

Proposition

Let X be a metric space. Let $u : X \rightarrow M$ be a map to a measure space (M, μ) . Then

$$\forall p \geq 1, \quad E_u^p(X) \leq \int_M E_u^{p-1}(u^{-1}(m)) d\mu(m).$$

$$\begin{aligned} \sum_i \mu(u(B_i))^p &= \sum_i \left(\int_M 1_{u(B_i)}(m) d\mu(m) \right) \mu(u(B_i))^{p-1} \\ &= \int_M \left(\sum_i 1_{u(B_i)}(m) \mu(u(B_i))^{p-1} \right) d\mu(m) \\ &= \int_M \left(\sum_{\{i; m \in u(B_i)\}} \mu(u(B_i))^{p-1} \right) d\mu(m) \\ &\leq \int_M E_u^{p-1; 2\epsilon} d\mu(m). \quad \text{q.e.d.} \end{aligned}$$

If u is Lipschitz and $\mu(B) \sim \text{radius}(B)^d$ in M , $E_u^{p-1} \leq \Pi^{d(p-1)}$, thus

$$E_u^p > 0 \quad \Rightarrow \quad \exists m, \dim(u^{-1}(m)) \geq d(p-1).$$

Corollary

Pick a submersion $v : \mathbb{R}^3 \rightarrow \mathbb{R}^d$. Assume that for all homeomorphisms $g : \text{Heis} \rightarrow \mathbb{R}^3$, $E_{v \circ g}^{4/d} > 0$. Then $\alpha_{qs}(G) \leq \frac{3-d}{4-d}$.

Corollary

Pick a submersion $v : \mathbb{R}^3 \rightarrow \mathbb{R}^d$. Assume that for all homeomorphisms $g : Heis \rightarrow \mathbb{R}^3$, $E_{v \circ g}^{4/d} > 0$. Then $\alpha_{qs}(G) \leq \frac{3-d}{4-d}$.

Proposition

If $u : G \rightarrow \mathbb{R}$ is continuous and non constant, $E_u^4 > 0$.

Corollary: $\alpha_{qs}(G) \leq \frac{2}{3}$.

Proof Length-area method. Let Γ denote the family of unit segments parallel to a side of unit n -cube. For all positive functions ρ on the square,

$$\int_0^1 \left(\int_{\gamma} \rho ds \right) d\gamma \leq \int \rho^n dx_1 \dots dx_n.$$

Replace integrals with packing (resp. covering) measures, apply to family of parallel horizontal line segments in $Heis$.

Proposition

Let X be a metric space. Let Γ be a family of subsets of X , equipped with a measure $d\gamma$. For each $\gamma \in \Gamma$, a probability measure m_γ is given on γ . Let $p \geq 1$. Assume that

$$\int_{\{\gamma \in \Gamma; \gamma \cap B \neq \emptyset\}} m_\gamma(\gamma \cap B)^{1-p} d\gamma \leq \tau.$$

Then, for every function ϕ on the set of balls of X ,

$$\Phi^p(X) \geq \frac{1}{\tau} \int_{\Gamma} \tilde{\Phi}^1(\gamma)^p d\gamma.$$

Proof Let $1_i(\gamma) = 1$ iff $\gamma \cap B_i \neq \emptyset$. The balls such that $1_i(\gamma) = 1$ cover γ , thus

$$\tilde{\Phi}^{1;\epsilon}(\gamma) \leq \sum_i \phi(B_i) 1_i(\gamma) = \sum_i \phi(B_i) 1_i(\gamma) m_\gamma(\gamma \cap B_i)^{\frac{1-p}{p}} m_\gamma(\gamma \cap B_i)^{\frac{p-1}{p}}.$$

Hölder's inequality gives

$$\tilde{\Phi}^{1;\epsilon}(\gamma)^p \leq \left(\sum_i \phi(B_i)^p 1_i(\gamma) m_\gamma(\gamma \cap B_i)^{1-p} \right) \left(\sum_i m_\gamma(\gamma \cap B_i) \right)^{p-1}.$$

Integrate over Γ .

Apply above Corollary to $d = 2$?

Unfortunately, there exist Lipschitz open maps $u : Heis \rightarrow \mathbb{R}^2$ such that $E_u^{4/3} < \infty$,
whence $E_u^2 = 0$.

Apply above Corollary to $d = 2$?

Unfortunately, there exist Lipschitz open maps $u : \text{Heis} \rightarrow \mathbb{R}^2$ such that $E_u^{4/3} < \infty$,
whence $E_u^2 = 0$.

Note that $E_u^2 = 0 \Rightarrow Du$ is not onto almost everywhere.

Question

Let $g : \text{Heis} \rightarrow \mathbb{R}^3$ be Lipschitz. Assume that the differential of g has rank ≤ 1 almost everywhere. Show that g cannot be a homeomorphism.